

# American Journal of Sports Medicine

<http://ajs.sagepub.com>

---

## **The Effect of Growth Factors on Biomechanical Properties of the Bone–Patellar Tendon–Bone Graft After Anterior Cruciate Ligament Reconstruction: A Canine Model Study**

Kazunori Yasuda, Fumihisa Tomita, Shuji Yamazaki, Akio Minami and Harukazu Tohyama  
*Am. J. Sports Med.* 2004; 32; 870 originally published online Apr 16, 2004;  
DOI: 10.1177/0363546503261695

The online version of this article can be found at:  
<http://ajs.sagepub.com/cgi/content/abstract/32/4/870>

---

Published by:

 SAGE Publications

<http://www.sagepublications.com>

On behalf of:



American Orthopaedic Society for Sports Medicine

**Additional services and information for *American Journal of Sports Medicine* can be found at:**

**Email Alerts:** <http://ajs.sagepub.com/cgi/alerts>

**Subscriptions:** <http://ajs.sagepub.com/subscriptions>

**Reprints:** <http://www.sagepub.com/journalsReprints.nav>

**Permissions:** <http://www.sagepub.com/journalsPermissions.nav>

**Citations** (this article cites 35 articles hosted on the  
SAGE Journals Online and HighWire Press platforms):  
<http://ajs.sagepub.com/cgi/content/abstract/32/4/870#BIBL>

# The Effect of Growth Factors on Biomechanical Properties of the Bone–Patellar Tendon–Bone Graft After Anterior Cruciate Ligament Reconstruction

## A Canine Model Study

Kazunori Yasuda,<sup>\*†‡</sup> MD, PhD, Fumihisa Tomita,<sup>‡</sup> MD, PhD, Shuji Yamazaki,<sup>‡</sup> MD, PhD, Akio Minami,<sup>‡</sup> MD, PhD, and Harukazu Tohyama,<sup>†</sup> MD, PhD

*From the Department of Sports Medicine and Joint Reconstruction Surgery, and <sup>‡</sup>the Department of Orthopaedic Surgery, Hokkaido University School of Medicine, Sapporo, Japan*

---

**Background:** No studies have dealt with the effect of growth factors on the free tendon autograft in anterior cruciate ligament reconstruction.

**Hypothesis:** Application of exogenous transforming growth factor- $\beta$  and epidermal growth factor may affect the structural properties and histology of the bone–patellar tendon–bone autograft after anterior cruciate ligament reconstruction.

**Study Design:** Controlled laboratory study.

**Methods:** Twenty dogs underwent anterior cruciate ligament reconstruction with the autogenous bone–patellar tendon–bone graft in bilateral knees. In 10 animals, 12 ng transforming growth factor- $\beta$  and 300 ng epidermal growth factor mixed with fibrin sealant of 0.6 mL were applied to the left knee. In the remaining 10 dogs, fibrin sealant alone was applied to the left knee. No additional treatments were applied to the right knee.

**Results:** The growth factor application increased the stiffness and maximum failure load of the femur-graft-tibia complex at 12 weeks ( $P = .016$  and  $P = .012$ , respectively); the sham treatment did not significantly affect them. Histologically, most of the cells in the grafts treated with growth factors had spindle-shaped nuclei; cells in the other grafts had round-shaped nuclei.

**Conclusions:** Application of transforming growth factor- $\beta$  and epidermal growth factor improves the structural properties of the autograft after anterior cruciate ligament reconstruction in the canine model.

**Clinical Relevance:** Application of growth factors is a possible strategy to prevent graft deterioration in anterior cruciate ligament reconstruction.

**Keywords:** anterior cruciate ligament (ACL) reconstruction; bone-tendon-bone autograft; transforming growth factor- $\beta$  (TGF- $\beta$ ); biomechanics; growth factor

---

\*Address correspondence to Kazunori Yasuda, MD, PhD, Department of Sports Medicine and Joint Reconstruction Surgery, Hokkaido University School of Medicine, Kita-ku, Kita-15, Nishi-7, Sapporo, 060-8638, Japan.

No author or related institution has received financial benefit from research in this study.

It has been established that structural properties of the femur-tendon autograft-tibia complex are reduced after ACL reconstruction and that the reduced properties are not restored even at 12 months after surgery.<sup>7,10</sup> The deterioration of the autograft is likely induced by extrinsic fibroblast proliferation and ingrowth along with revascularization, which occur after intrinsic fibroblast necrosis.<sup>19,33</sup> Therefore, one future goal in ACL reconstruction is

to prevent the graft deterioration after transplantation, or to accelerate mechanical restoration of the deteriorated graft. Recently, a number of *in vitro* studies have shown that application of various growth factors significantly affects fibroblasts harvested from tendon and ligament tissues. For example, transforming growth factor- $\beta$  (TGF- $\beta$ ) increases both collagen and noncollagenous protein synthesis by fibroblasts<sup>11,23,25,27</sup> and stimulates fibroblast proliferation.<sup>20,30,31</sup> Epidermal growth factor (EGF) also stimulates fibroblast proliferation *in vitro*.<sup>11,30</sup> Combined application of these 2 growth factors enhances their effects.<sup>4,28,32</sup> In addition, several *in vivo* studies have reported that some growth factors significantly enhance healing of the injured medial collateral ligament.<sup>7,14,21,37,40</sup> Therefore, there is the possibility that intra-articular administration of these growth factors may affect remodeling of the graft in ACL reconstruction.


Recently, the authors<sup>5,29</sup> have studied the effect of TGF- $\beta$  and EGF on the *in situ* frozen-thawed rabbit ACL, an idealized model of anatomic but acellular autograft. We have found that intra-articular administration of TGF- $\beta$  and EGF not only significantly inhibits reduction of the material properties of the frozen-thawed ACL but also increases the cross-sectional area, and the administration changes the diameter distribution of collagen fibrils in the frozen-thawed ACL. However, this model is not completely equivalent to an ACL reconstruction model with the free tendon graft. No studies have dealt with the effect of growth factors on the free tendon autograft in ACL reconstruction.

Based on the above-mentioned studies,<sup>5,29</sup> we have hypothesized that an intra-articular application of TGF- $\beta$  and EGF mixed with the fibrin sealant may significantly affect the structural properties and histology of the bone-patellar tendon-bone (B-PT-B) autograft in ACL reconstruction. The purpose of this study is to test this hypothesis with a canine model.

## MATERIALS AND METHODS

### Study Design

A biomechanical study was conducted with 25 healthy, adult beagle dogs weighing  $10.9 \pm 0.6$  kg (mean  $\pm$  SD). Animal experimentation was carried out under the Role and Regulation of the Animal Care and Use Committee, Hokkaido University, School of Medicine. Twenty dogs underwent ACL reconstruction with the autogenous B-PT-B graft in bilateral knees (Figure 1). Then, in 10 of the 20 animals, 12 ng TGF- $\beta$  and 300 ng EGF (R&D System, Minneapolis, Minn) mixed with fibrin sealant of 0.6 mL (Kaketsuken Co, Kumamoto, Japan) were applied around the graft in the left knee (GF group), and fibrin sealant alone was applied as the sham treatment in the right knee (sham group). In the remaining 10 dogs, no additional treatments were applied after surgery, and only the right knee was used to obtain control data of natural ACL reconstruction with the B-PT-B autograft (control group). No animals were immobilized postoperatively, and all were allowed unrestricted activities in their cages (1 dog per



**Figure 1.** The ACL reconstruction procedures with the bone-patellar tendon-bone graft.

cage). In addition, bilateral knees harvested from the remaining 5 dogs, which underwent no surgery, were used to obtain normal biomechanical data of the ACL. All animals were sacrificed by a lethal injection of thiamylal sodium at 12 weeks after the operation. In each group ( $n = 10$ ), 8 and 2 knees were used for biomechanical and histological examinations, respectively.

### Growth Factors and a Delivery Vehicle

Recombinant human TGF- $\beta$ 1 and recombinant human EGF were purchased from R&D Systems. Commercially available fibrin sealant (Kaketsuken Co) was used as a vehicle for the growth factors.<sup>14,40</sup> The fibrin sealant was formed by human fibrinogen (80 mg/mL) and thrombin being mixed in the presence of calcium and factor XIII. The TGF- $\beta$ 1 and EGF were reconstituted according to the instructions given by the manufacturer. The reconstituted growth factors were mixed with the thrombin solution, and then the fibrinogen-thrombin solution was combined. The resultant fibrin sealant of 0.6 mL containing growth factors was applied to each knee.

### Surgical Procedure and Postoperative Treatment

Each procedure was performed under general endotracheal anesthesia. Using sterile techniques, a medial parapatellar arthrotomy was performed in each knee. The anterior aspect of the infrapatellar fat pad was excised to allow

for a full visualization of the tibial attachment of the ACL. The ACL in both knees was excised.

The B-PT-B preparation having a width of 4 mm and a bone plug length of 10 mm was harvested from the right extensor apparatus of the knee, and the bone plugs were then trimmed so that the plug could be passed through a 4-mm-diameter sheath. The periosteum, which covered the anterior (ventral) aspect of the bone plug, was not dissected. A drill hole having a diameter of 1 mm was made in each bone plug, and a No. 1 nonabsorbable polyester suture (Ticron, Davis and Geck, Wayne, NJ) was passed through each hole. After the preparation, the cross-sectional area of the middle portion of each graft was measured with an area micrometer, which had been established in our previous studies.<sup>22,26</sup> In brief, the middle portion was located in the micrometer slot, and the plunger was placed on the specimen in the slot. The thickness of the specimen was measured while it was being pressed under a constant pressure of 0.12 MPa. The cross-sectional area of the middle portion was calculated by multiplying the slot width by the measured thickness.

The anteromedial surface of the tibia was exposed, and the periosteum was elevated. A bone tunnel having a diameter of 4 mm was drilled in the tibia through the central tibial insertion of the resected ACL to the exposed anteromedial surface of the tibia. Then, a small incision was made in the posterolateral part of the femoral joint capsule. A sharp curette was inserted from the incision to over the top of the lateral femoral condyle, and the cortical bone at this portion was then curetted along the over-the-top route so that a "trough" was created. For each graft, the distal end having a length of 15 mm was placed in the tibial bone tunnel, and the proximal end was placed in the trough. This over-the-top trough technique for the femoral side was performed because it had been used as a reliable and standard technique for the canine knee,<sup>3,34</sup> and the additional trough technique enabled this technique to resemble the tunnel technique for human patients. The tibial bone plug was grafted so that the periosteum surface faced the anterior wall of the tunnel. Consequently, the opposite surface of the bone plug faced the posterior wall. We tethered each end of the graft with the suture to a screw inserted into the bone, manually applying minimal graft tension in a direction aligned with the long axis of the bone tunnel with the knee flexed at 45° (Figure 1). After graft fixation, we found no anteroposterior instability by manual tests. The surgical wound was irrigated with a physiologic saline solution and closed with 3-0 nylon sutures.

None of the beagles were immobilized postoperatively. (70 cm in width, 68 cm in height, and 70 cm in depth). In this cage, each beagle could walk along the cage wall, although it could not run. At the time of sacrifice, first, anesthesia was induced by the intramuscular administration of ketamine hydrochloride (10 mg/kg). Then, a lethal dose of thiethylal sodium was injected intravenously.

### Sample Preparation for Mechanical Testing

Each hindlimb for mechanical testing was wrapped in gauze moistened with physiologic saline solution and then

wrapped in air-tight plastic films. The specimens were stored at -32°C until testing. All mechanical tests were performed blinded; namely, the researcher who performed the tests was different from the surgeon for this experimental operation, and the researcher was blinded concerning the group name of each specimen at the time of testing. Before mechanical testing, each knee was thawed overnight at 4°C. The knee was removed from the hindlimb, and the surrounding muscles were carefully removed.

### Biomechanical Testing

The knee specimen with the femur having a length of 45 mm and the tibia having a length of 60 mm was removed from the animal immediately after sacrifice. Each specimen for biomechanical testing was wrapped in gauze moistened with physiologic saline solution and then wrapped in an air-tight polychlorovinylidene film. The specimens were stored at -32°C until the time of testing. Prior to mechanical testing, each knee was thawed overnight at 4°C. The anterior-posterior translation was evaluated according to Beynon et al.<sup>6</sup> The specimen was mounted on the fixture of 5 degrees of freedom at 45° and 90° of knee flexion, and 50-N anterior and posterior shear loads were applied to the knee specimen.<sup>17,18</sup>

All soft tissues other than the graft were carefully dissected. Then the cross-sectional area of the graft was measured with the above-described area micrometer. The femur and the tibia were separately cast in rectangular aluminum tubes using polymethylmethacrylate resin.

With the use of a set of specially designed grips, the prepared femur-graft-tibia complex specimen was attached to a conventional tensile tester (PMT 250W; Orientec, Tokyo, Japan) so that the tibia was positioned to allow for tensile loading aligned with the long axis of the bone tunnel with the knee flexed at 45° (Figure 2). Two parallel lines were drawn transversely on the proximal bone surface of the femur and the tibia using Nigrosine stain as gauge-length markers to measure elongation of the femur-graft-tibia complex. Prior to the tensile test of the femur-graft-tibia complex, the specimen was preconditioned with a static preload of 0.5 N for 5 minutes, followed by 10 cycles of loading and unloading with a strain of 0.5% at a crosshead speed of 20 mm per minute.<sup>17,18</sup> Then, the femur-graft-tibia complex specimen was stretched until failure at the same rate, and the load-deformation curve was drawn on a recorder (X-Y-T Recorder 3023; Yokogawa, Tokyo, Japan). The elongation of the complex was determined with a video dimension analyzer (HTV-C1170; Hamamatsu Photonics, Tokyo, Japan) using the previously described gauge-length markers.<sup>26</sup> The specimen was kept moistened throughout the test period with a physiological saline solution spray. During tensile testing, failure modes were carefully observed in each femur-graft-tibia complex specimen. From the load-elongation curve, the linear stiffness was determined as the slope of the linear portion. In addition, the maximum failure load and the elongation at failure of the femur-graft-tibia complex were obtained on the load-elongation curve.




Figure 2. Tensile testing for the femur-graft-tibia complex.

Histological Observation

The graft-tibia specimen for histological observation was fixed in a 10% buffered formalin solution immediately after harvesting from each limb. After the specimen was decalcified, it was cast in paraffin blocks. Each specimen was sectioned parallel to the longitudinal axis of both the tendon graft and the bone plug, stained with hematoxylin and eosin and toluidine blue for light and polarized light microscopy.

Statistical Analysis

All data were described with the mean and SD values. Each biomechanical parameter was compared with the 1-way analysis of variance. The chi-square test was used to compare the failure mode between the groups. The most essential comparison was made between the GF and sham groups to test the above-described hypothesis. To support this comparison, additional comparisons were performed between the sham and control groups and between the GF group and the normal knee. The significance level was set at  $P = .05$  for each test. The statistical software (Stat View, Abacus Ontent Inc, Berkeley, Calif) was used for the analysis.

RESULTS

Gross Observations

Postoperatively, the animals carried the involved limbs for about a week. Their cage activities gradually increased,




Figure 3. The anterior-posterior (A-P) translation of the tibia on the femur measured at 45° and 90° of flexion. There were no significant differences between the graft (GF) group and the normal knee, although the sham and control groups were significantly greater than the normal knees. There were no significant differences between the GF and sham groups.

TABLE 1  
The Anterior-Posterior Translation of the Tibia at 12 Weeks (Mean ± SD)

Knee Flexion Angle	Group			
	Graft	Sham	Control	Normal
45°	4.7 ± 1.1	5.3 ± 1.4	6.8 ± 1.7	3.5 ± 0.4
90°	4.7 ± 1.3	5.8 ± 1.7	5.6 ± 1.1	2.8 ± 0.4

and by the fourth postoperative week, the animals showed only a mild degree of lameness. No lameness was noted after the sixth week. At the time of sacrifice, no graft failure was found in any of the knees. The intra-articular graft was enveloped by granulation tissues, which adhered to the infrapatellar fat pad, in each group. The adhesion appeared to an obviously greater degree in the GF and sham groups than in the control group. The synovial membrane was also thickened around the patellofemoral joint to an obviously greater degree in the GF and sham groups than in the control group. Neither meniscus tears nor tibiofemoral osteophyte formation was found in each knee. In addition, a small part of the surface of articular cartilage showed mild degeneration, such as softening and fibrillation, in each knee.


Anterior-Posterior Translation of the Tibia

Concerning the anterior-posterior translation of the tibia on the femur measured at 45° of flexion, there were no significant differences between the GF and sham groups (Figure 3). However, the sham group (average 5.3 mm) as well as the control group (6.8 mm) were significantly greater ( $P = .012$  and  $P = .001$ , respectively) than the nor-



TABLE 2  
The Cross-sectional Area and the Structural Properties of the Femur-Graft-Tibia Complex Measured at 12 Weeks (Mean  $\pm$  SD)

Femur-Graft-Tibia Complex	Group			
	Graft	Sham	Control	Normal
Cross-sectional area (mm <sup>2</sup> )	12.7 $\pm$ 2.1	9.6 $\pm$ 2.0	10.8 $\pm$ 3.6	9.8 $\pm$ 1.6
Stiffness (N/mm)	93.8 $\pm$ 20.1	62.1 $\pm$ 32.1	54.4 $\pm$ 17.8	130.4 $\pm$ 35.6
Maximum load (N)	303.2 $\pm$ 107.5	184.4 $\pm$ 90.4	176.4 $\pm$ 74.1	626.0 $\pm$ 108.5
Elongation at failure (mm)	4.7 $\pm$ 1.0	4.7 $\pm$ 0.8	4.9 $\pm$ 1.4	7.0 $\pm$ 2.1



**Figure 4.** The cross-sectional area of the tendon graft measured at 12 weeks. The area was significantly greater in the graft (GF) group than in the sham group, whereas there were no significant differences between the sham and control groups. Note that the area measured immediately before implantation averaged  $4.9 \pm 0.5$  mm<sup>2</sup>.


mal knees (3.5 mm), whereas there were no significant differences between the GF group (4.7 mm) and the normal knee. Concerning the translation measured at 90° of flexion, we found the same tendency as observed at 45°, except that the GF group was also significantly greater than the normal knees (Table 1).

#### Cross-sectional Area of the Graft

The cross-sectional area of the tendon graft measured immediately before implantation averaged  $4.9$  mm<sup>2</sup> with a SD of 0.5, which was approximately half that of the normal ACLs (average  $9.9$  mm<sup>2</sup>, SD  $\pm 1.2$ ). In each group, the cross-sectional area significantly increased twice or more at 12 weeks, compared to the preoperative value. The area was significantly greater ( $P = .017$ ) in the GF group ( $12.7$  mm<sup>2</sup>) than in the sham group ( $9.7$  mm<sup>2</sup>) at 12 weeks, whereas there were no significant differences between the sham and control groups (Figure 4). In addition, the cross-

TABLE 3  
Failure Modes in Tensile Tests

Mode	Group			
	Graft	Sham	Control	Normal
Midsubstance tear	4	8	8	0
Avulsion fracture	4	0	0	10




**Figure 5.** The load-elongation curves of the femur-graft-tibia complex in graft (GF), sham, and control groups and the normal knee.


sectional area of the GF group was significantly greater ( $P = .026$ ) than that of the normal ACLs (Table 2).

#### Structural Properties of the Femur-Graft-Tibia Complex

Concerning failure modes of the femur-graft-tibia complex, the tendon graft portion was torn at the midsubstance in 4 knees and avulsed from the tibia with a small fragment in the remaining 4 knees in the GF group. The remaining part of the bone plug that had been grafted within the tunnel was not identified from the tibia. In the sham group, the tendon graft portion was torn at the midsubstance in each knee. There were significant differences in the failure



**Figure 6.** The stiffness of the femur-graft-tibia complex measured at 12 weeks. The graft (GF) group was significantly greater than the sham group, whereas it was significantly lower than the normal knees. There were no significant differences between the sham and control groups.



**Figure 7.** The maximum failure load of the femur-graft-tibia complex measured at 12 weeks. The graft (GF) group was significantly greater than the sham group, whereas it was significantly lower than the normal knees. There were no significant differences between the sham and control groups.

mode between the 2 groups (chi-square test,  $P = .025$ ). In the control group, the graft portion was torn at the mid-substance in each knee, whereas the normal ACL was avulsed from the tibia with a small fragment in each knee (Table 3). The size of the avulsed fragment was similar between the GF group and the control group.

Figure 5 shows the load-elongation curves of the femur-graft-tibia complex in the GF, sham, and control groups and the normal knee. The load-elongation curve of the GF group was obviously different from not only the curves of

the sham and control groups but also the curve of the normal knee.

The stiffness of the femur-graft-tibia complex in the GF group (average, 93.8 N/mm) was significantly greater ( $P = .016$ ) than that in the sham group (62.1 N/mm) at 12 weeks, whereas it was significantly lower ( $P = .013$ ) than that of the normal knees (130.4 N/mm). There were no significant differences between the sham and control groups (Figure 6).

Concerning the maximum failure load of the femur-graft-tibia complex, the GF group (average, 303.2 N) was significantly greater ( $P = .012$ ) than the sham group (184.4 N) at 12 weeks, whereas it was significantly lower ( $P < .0001$ ) than that of the normal knees (626.0 N). There were no significant differences between the sham and control groups (Figure 7).

Concerning the elongation at failure of the femur-graft-tibia complex, there were no significant differences between the GF, sham, and control groups. Each group was significantly lower than the normal knees ( $P = .003$ ,  $P = .003$ , and  $P = .007$ , respectively) (Table 2).


#### Histologic Observation of the Graft

The collagen fibers in the GF group appeared to be more oriented compared to those in the control and sham groups (Figures 8 A, B, and C). The latter fibers appeared to be coarse (Figures 8 B and C). In the control and sham groups, numerous cells that had a round-shaped nucleus were randomly scattered in the midsubstance of the graft (Figure 8 B and C). In the GF group (Figure 8A), almost all of the cells had a spindle-shaped or oval-shaped nucleus. In this group, the number of cells scattering in the midsubstance appeared to be less than that in the control and sham groups. In the GF group, in addition, we observed several cells with an oval-shaped nucleus that were longitudinally aligned between collagen fascicles, parallel to the fibers (Figure 8A). However, the histology involving cells and collagen fibers of the GF group was different from that of the normal ACL (Figure 8D).

Observations using transillumination by polarized light showed that a regular crimp pattern of collagen fibers with relatively large amplitude and wavelength was predominantly observed in the GF group (Figure 9A). In the control and sham groups, a crimp pattern was irregular (Figures 9 B and C). However, the crimp pattern of the GF group was different from that of the normal ACL, which showed regular crimp with amplitude and wavelength less than those of the GF group (Figure D).

#### DISCUSSION

This study clearly demonstrated that intra-articular administration of 12 ng TGF- $\beta$  and 300 ng EGF mixed with fibrin sealant significantly increases the cross-sectional area of the tendon graft, significantly changes the failure mode of the femur-graft-tibia complex, and significantly increases the stiffness of the femur-graft-tibia complex in the canine ACL reconstruction model. Histologically, this



**Figure 8.** Histology of the midsubstance. The collagen fibers in the graft (GF) group (A) appeared to be more oriented compared to those in the control group (B) and the sham group (C). The latter fibers appeared to be coarse. Note the number, shape, and alignment of cells of the GF group compared to the control and the sham groups (see text). However, the histology of the GF group was different from that of the normal ACL (D).


study implied that the administration of these growth factors affects cell maturation in the tendon graft. Thus, this is the first article that reported that the intra-articular administration of TGF- $\beta$  and EGF significantly affects the structural properties of the femur-autograft-tibia complex reconstructed in ACL reconstruction with the B-PT-B graft.

A number of studies have shown the natural course of the tendon autograft after ACL reconstruction.<sup>1-3,9,10,19</sup> Following transplantation, intrinsic fibroblasts in the tendon graft undergo a process of ischemic necrosis. The graft is then enveloped with a synovial tissue by 2 weeks, and extrinsic cell infiltration with revascularization occurs in the graft. Butler et al<sup>7</sup> and Clancy et al<sup>10</sup> reported that the cross-sectional area significantly increases at approximately 6 weeks after ACL reconstruction surgery and that the structural properties of the femur-graft-tibia complex dramatically decrease throughout the observation period, compared to the normal opposite knee. Ultrastructurally, the population of small-diameter fibrils increases in the frozen-thawed goat ACL at 6 months.<sup>13,16</sup> In the present

study, the histological and biomechanical results observed in the control group at 12 weeks were similar to the natural course of the patellar tendon graft reported in the above-described studies. Therefore, we can say that the control data in this study were not unusual, compared to the above-described studies with a canine model.

Concerning the effect of fibrin sealant, previous studies have reported that the application of fibrin sealant did not give any significant effects on remodeling of the frozen-thawed ACL<sup>5,29</sup> or the injured ligament.<sup>14,40</sup> The present study also showed that application of only fibrin sealant had no significant effects on structural properties of the femur-graft-tibia complex and histology of the graft. This fact also showed that the data in the sham group were not unusual in comparison with the above-described studies. However, the adhesion between the graft and the infrapatellar fat pad appeared to an obviously greater degree in the GF and sham groups than in the control group. The synovial membrane was also thickened around the patellofemoral joint to an obviously greater degree in the GF and sham groups than in the control group. These find-





**Figure 9.** Observations using transillumination by polarized light. In the graft (GF) group (A), a regular crimp pattern of collagen fibers with relatively large amplitude and wavelength was predominantly observed. In the control group (B) and the sham group (C), a crimp pattern was irregular. However, the crimp pattern of the GF group was different from that of the normal ACL (D).

ings imply that the fibrin sealant, which was formed by human fibrinogen and thrombin being mixed, gave some adverse effects to the fat and synovial tissues, although it did not significantly affect the biomechanical data. In addition, we should recognize the possibility that TGF- $\beta$  and EGF may have detrimental effects on the tissues within the joint. For example, Van Beuningen et al<sup>36</sup> and Hulth et al<sup>15</sup> reported that repetitive application of TGF- $\beta$  into the knee joint induces degenerative changes in the cartilage. In this study, however, we applied these growth factors only once after surgery. This may be the reason we did not find any significant detrimental effects in this study.

Based on these fundamental results, the present study showed that the intra-articular application of TGF- $\beta$  and EGF with fibrin sealant significantly affects the biomechanical properties of the femur-graft-tibia complex after ACL reconstruction. For example, the graft was avulsed from the tibia with a small fragment in 4 of the 8 knees in the GF group during tensile tests, whereas each complex failed at the midsubstance of the graft in the sham and control groups. The avulsion from the tibia with a small fragment is a common failure mode in the normal ACL.<sup>38,39</sup>

In this study, the size of the avulsed fragment was similar between the GF group and the control group, and the remaining part of the bone plug that had been grafted within the tunnel was not identified from the tibia. In this study, therefore, we estimated the bone plug to be healed within the tunnel. Our previous study with the same canine model showed that the bone plug was healed within the tibial tunnel at 6 weeks after surgery.<sup>34</sup> This study supports our estimation on bone plug healing. In addition, the maximum failure load of the femur-graft-tibia complex of the GF group was significantly greater than that of the sham group. It was noted that the growth factor application significantly increased the stiffness of the femur-graft-tibia complex, compared to the sham and control groups. This result was considered to show the stiffness of the graft substance to a greater degree than that of the insertion of the graft. Thus, these results suggested that the application of TGF- $\beta$  and EGF significantly suppresses the natural deterioration of the structural properties of the tendon autograft in ACL reconstruction.

We should speculate on a few possible mechanisms to explain why the structural properties of the graft were

changed by the application of TGF- $\beta$  and EGF, although we did not definitely determine them in this study. First, in the present study, the administration of TGF- $\beta$  and EGF significantly increased the cross-sectional area of the tendon graft. Marui et al<sup>23</sup> reported that TGF- $\beta$  increases not only collagen synthesis but also noncollagenous protein synthesis in ligament fibroblasts. Therefore, there is the possibility that in the present study, an application of the growth factors enhanced synthesis of new tissues involving various collagens, bonding proteins, and/or proteoglycans in the graft. These changes might affect the proteoglycan-collagen network and/or minor collagen network in the graft. An increase of the cross-sectional area does not always mean an increase of the structural properties. In this study, however, for example, if causes of the increase of the cross-sectional area involved enhanced production of type III and/or type I collagens in comparison with the natural course, it is considered that it was one of the causes of the increase of the structural properties. Second, histological observations in this study implied that the administration of these growth factors affects cell maturation in the tendon graft. We should note in the natural course of the graft that the structural properties of the graft are extremely deteriorated after surgery. A number of previous studies have reported that TGF- $\beta$  suppresses collagenase activities from fibroblasts in various tissues.<sup>24,35</sup> In the present study, therefore, there is the possibility that an application of the growth factors suppressed the collagenase activity from extrinsic fibroblasts infiltrating in the graft, resulting in the inhibition of their mechanical deterioration. These mechanisms can simultaneously occur. Further studies should be conducted to clarify the mechanisms or causes of the increase of the structural properties of the graft.

Concerning the anterior-posterior translation of the knee, all the operated knees appeared to be looser than the normal knee. It has been well known that this phenomenon commonly occurs in animal models.<sup>7,10,16</sup> It shows that the graft is elongated to some degree in the animal model. Therefore, we should recognize that the natural course of the graft is not completely the same as that in human patients. However, we also note that results from such animal models have contributed to the progression of the ACL reconstruction. In the present study, although the sham and control groups were significantly greater than the normal knees, there were no significant differences between the GF group and the normal knee. Therefore, although there were no significant differences between the GF and sham groups, some tendencies exist that indicate the application of TGF- $\beta$  and EGF improved the translation. However, we could not come to a conclusion about the effect of the application of TGF- $\beta$  and EGF on the length of the graft.

It is necessary to recognize some limitations concerning the experimental model used in this study. First, we used the beagle model in this study. The average maximum load of the normal ACL was 626 N. This value is higher than the value of approximately 400 N shown in previous studies with beagles.<sup>12</sup> Therefore, we believe that our testing

condition was acceptable. However, this value was less than the maximum load of 1210 N reported with German shepherd dogs.<sup>8</sup> Among various canine models, this model is recommended for use in experimental studies from an ethical viewpoint. However, we should recognize that the beagle model is relatively small in various canine models. Second, we used the bilateral injury model. In this model, it is considered that animals have to evenly load the graft in each knee during walking. This is an advantage of this model but simultaneously is a disadvantage. Namely, there is a possibility that the graft might be overloaded in the early phase after ACL reconstruction and that the over-stress might give detrimental effects to the graft, such as an increase of the anterior-posterior translation of the knee.<sup>17</sup> However, the bilateral injury model enabled us to compare the 2 groups under the same postoperative environment concerning the effect of the growth factor application, which was the main purpose of this study.

There are some additional limitations to this study. The first limitation was that we could not determine the structural and mechanical properties of the graft itself, although we dealt with the whole femur-graft-tibia complex. However, the structural properties of the whole femur-graft-tibia complex are the most essential factors to evaluate the knee with ACL reconstruction from the clinical viewpoint, because not only the graft midsubstance but also insertion of the graft are equally important for success of ACL reconstruction. The second limitation was that we did not evaluate long-term results after ACL reconstruction. However, we believe that this study is of value because it first demonstrated that the application of TGF- $\beta$  and EGF significantly affects the biomechanical properties of the free tendon graft in ACL reconstruction. The third limitation was that we did not examine the diameter of collagen fibrils with a translucent electron microscope. The fourth limitation is a potential bias in the histological observations in this study. The sample number was only 2 in each group. In addition, our observation was traditionally performed according to the previous studies.<sup>1,3,9,10,16</sup> Therefore, we did not perform random sampling or observation in a blinded fashion, nor did we carry out statistical analyses on histological observations. Based on the biomechanical data with the statistical significance, we should deal with the histological findings as additional data supporting the biomechanical data. Further studies should be conducted with a sufficient number of more appropriate animals to answer questions concerning each limitation.

Finally, it has been recognized that there are various biological and biomechanical factors affecting the outcome of ACL reconstruction.<sup>41</sup> This study also suggested that growth factors are important factors affecting results of ACL reconstruction. As to the clinical relevance of this study, this study implied that the application of certain growth factors is of value to be studied as one of the strategies for future ACL reconstruction to prevent graft deterioration after transplantation or to accelerate mechanical restoration of the deteriorated graft. However, we also should recognize the possibility that growth factors may

have some detrimental effects on the tissues within the joint, such as detrimental effects of TGF- $\beta$  on the joint cartilage. Therefore, in the near future, we should study how to effectively apply such growth factors to the targeted tissue without their detrimental effects.

## ACKNOWLEDGMENT

This work was supported financially in part by the Grant-in-Aid for Scientific Research (12470299 and 14GS0301) from the Ministry of Education, Science and Culture, Japan.

## REFERENCES

- Alm A, Ekstrom H, Gillquist J, Stromberg B. The anterior cruciate ligament: a clinical and experimental study on tensile strength, morphology and replacement by patellar ligament. *Acta Chir Scand*. 1974;445(suppl):4-49.
- Amiel D, Kleiner JB, Akeson WH. The natural history of the anterior cruciate ligament autograft of patellar tendon origin. *Am J Sports Med*. 1986;14:449-462.
- Arnoczky SP, Tarvin GB, Marshall JL. Anterior cruciate ligament replacement using patellar tendon: an evaluation of graft revascularization in the dog. *J Bone Joint Surg Am*. 1982;64:217-224.
- Assoian RK, Frolik CA, Roberts AB, et al. Transforming growth factor- $\beta$  controls receptor levels for epidermal growth factor in NRK fibroblasts. *Cell*. 1984;36:35-41.
- Azuma H, Yasuda K, Tohyama H, et al. Timing of administration of transforming growth factor- $\beta$  and epidermal growth factor influences the effect on material properties of the in situ frozen-thawed anterior cruciate ligament. *J Biomech*. 2003;36:373-381.
- Beynonn BD, Johnson RJ, Tohyama H, Renstrom PA, Arms SW, Fischer RA. The relationship between anterior-posterior knee laxity and the structural properties of the patellar tendon graft: a study in canines. *Am J Sports Med*. 1994;22:812-820.
- Butler DL, Grood ES, Noyes FR, et al. Mechanical properties of primate vascularized versus nonvascularized patellar tendon grafts. *J Orthop Res*. 1989;7:68-79.
- Cabaud HE, Rodkey WG, Feagin JA. Experimental studies of acute anterior cruciate ligament injury and repair. *Am J Sports Med*. 1979;7:18-22.
- Chiroff RT. Experimental replacement of the anterior cruciate ligament: a histological and microradiographic study. *J Bone Joint Surg Am*. 1975;57:1124-1127.
- Clancy WG, Narechania RG, Rosenberg TD, et al. Anterior and posterior cruciate ligament reconstruction in rhesus monkeys. *J Bone Joint Surg Am*. 1981;63:1270-1284.
- DesRosiers EA, Yahia L, Rivard CH. Proliferative and matrix synthesis response of canine anterior cruciate ligament fibroblasts submitted to combined growth factors. *J Orthop Res*. 1996;14:200-208.
- Foggie HE III, Bahniuk EH, Helpe KG, Davy DT. The effects of tibial-femoral angle on the failure mechanics of the canine anterior cruciate ligament. *J Biomech*. 1986;19:89-91.
- Frogameni AD, Jackson DW, Simon TM. Collagen remodeling in ACL reconstruction: goat model. In: Jackson DW, ed. *The Anterior Cruciate Ligament: Current and Future Concepts*. New York: Raven Press; 1993:219-226.
- Hildebrand KA, Woo SL-Y, Smith DW, et al. The effects of platelet-derived growth factor-BB on healing of the rabbit medial collateral ligament. *Am J Sports Med*. 1998;26:549-554.
- Hulth A, Johnell O, Miyazono K, Lindberg L, Heinegard D, Heldin C-H. Effect of transforming growth factor- $\beta$  and platelet-derived growth factor-BB on articular cartilage in rats. *J Orthop Res*. 1996;14:547-553.
- Jackson DW, Grood ES, Goldstein J, et al. A comparison of patellar tendon autograft and allograft used for anterior cruciate ligament reconstruction in goats. *Am J Sports Med*. 1993;21:176-185.
- Katsuragi R, Yasuda K, Tsujino J, et al. The effect of nonphysiologically high initial tension on the mechanical properties of *in situ* frozen anterior cruciate ligament in a canine model. *Am J Sports Med*. 2000;28:47-56.
- Keira M, Yasuda K, Kaneda K, Yamamoto N, Hayashi K. Mechanical properties of the anterior cruciate ligament chronically relaxed by elevation of the tibial insertion. *J Orthop Res*. 1996;14:157-166.
- Kleiner JB, Amiel D, Harwood FL, Akeson WH. Early histologic, metabolic, and vascular assessment of anterior cruciate ligament autografts. *J Orthop Res*. 1989;7:235-242.
- Lee J, Green MH, Amiel D. Synergistic effect of growth factors on cell outgrowth from explants of rabbit anterior cruciate and medial collateral ligaments. *J Orthop Res*. 1995;13:435-441.
- Letson AK, Dahners LE. The effect of combinations of growth factors on ligament healing. *Clin Orthop*. 1994;308:207-212.
- Majima T, Yasuda K, Yamamoto N, et al. Deterioration of mechanical properties of the autograft in controlled stress-shielded augmentation procedures. *Am J Sports Med*. 1994;22:821-829.
- Marui T, Niyibizi C, Georgescu HI, et al. Effects of growth factors on matrix synthesis by ligament fibroblasts. *J Orthop Res*. 1997;15:18-23.
- Mauviel A. Cytokine regulation of metalloproteinase gene expression. *J Cell Biochem*. 1993;53:288-295.
- Murphy PG, Loitz BJ, Frank CB, Hart DA. Influence of exogenous growth factors on the synthesis and secretion of collagen types I and III by explants of normal and healing rabbit ligaments. *Biochem Cell Biol*. 1994;72:403-409.
- Ohno K, Yasuda K, Yamamoto N, et al. Effects of complete stress-shielding on the mechanical properties and histology of in situ frozen patellar tendon. *J Orthop Res*. 1993;11:592-602.
- Pierce GF, Mustoe TA, Lingelbach J, et al. Platelet-derived growth factor and transforming growth factor- $\beta$  enhance tissue repair activities by unique mechanisms. *J Cell Biol*. 2000;109:429-440.
- Roberts AB, Anzano MA, Lamb LC, et al. Isolation from murine sarcoma cells of novel transforming growth factors potentiated by EGF. *Nature*. 1982;295:417-419.
- Sakai T, Yasuda K, Tohyama H, et al. Effects of combined administration of transforming growth factor- $\beta$  1 and epidermal growth factor on properties of the *in situ* frozen anterior cruciate ligament in rabbits. *J Orthop Res*. 2002;20:1345-1351.
- Schmidt CC, Georgescu HI, Kwok CK, et al. Effect of growth factors on the proliferation of fibroblasts from the medial collateral and anterior cruciate ligaments. *J Orthop Res*. 1995;13:184-190.
- Spindler KP, Imro AK, Mayes CE, Davidson JM. Patellar tendon and anterior cruciate ligament have different mitogenic responses to platelet-derived growth factor and transforming growth factor  $\beta$ . *J Orthop Res*. 1996;14:542-546.
- Sporn MB, Roberts AB, Shull JH, et al. Polypeptide transforming growth factors isolated from bovine sources and used for wound healing in vivo. *Science*. 1983;219:1329-1331.
- Tohyama H, Yasuda K. Extrinsic cell infiltration and revascularization accelerate mechanical deterioration of the patellar tendon after fibroblast necrosis. *J Biomech Eng*. 2000;122:594-599.
- Tomita F, Yasuda K, Mikami S, Sakai T, Yamazaki S, Tohyama H. Comparisons of intraosseous graft healing between the doubled flexor tendon graft and the bone-patellar tendon-bone graft in anterior cruciate ligament reconstruction. *Arthroscopy*. 2001;17:461-476.
- Uria JA, Jimenez MG, Balbin M, et al. Differential effects of transforming growth factor- $\beta$  on the expression of collagenase-1 and collagenase-3 in human fibroblasts. *J Biol Chem*. 1998;273:9769-9777.
- Van Beuningen HM, van der Kraan PM, Arntz OJ, van den Berg WB. Transforming growth factor- $\beta$  1 stimulates articular chondrocyte proteoglycan synthesis and induces osteophyte formation in the murine knee joint. *Lab Invest*. 1994;71:279-290.
- Weiss JA, Beck CL, Levine RE, Greenwald RM. Effects of platelet-derived growth factor on early medial collateral ligament healing. *Trans Orthop Res Soc*. 1995;20:159.



38. Woo SL-Y, Hollis JM, Adams DJ, et al. Tensile properties of the human femur–anterior cruciate ligament–tibia complex: the effects of specimen age and orientation. *Am J Sports Med.* 1991;19:217-225.
39. Woo SL-Y, Hollis JM, Roux RD, et al. Effects of knee flexion on the structural properties of the rabbit femur–anterior cruciate ligament–tibia complex. *J Biomech.* 1987;20:557-563.
40. Woo SL-Y, Taylor BJ, Schmidt CC, et al. The effect of dose levels of growth factors on the healing of the rabbit medial collateral ligament. *Trans Orthop Res Soc.* 1996;21:97.
41. Yasuda K, Tsujino J, Tanabe Y, Kaneda K. Effects of initial graft tension on clinical outcome after anterior cruciate ligament reconstruction. *Am J Sports Med.* 1997;25:99-106.